## 4 Phases of Healing **ROADMAP** During a Disc Episode

1. PAIN relief	2 CWELLING reduction	3. DISC DAMAGE	
happens in approximately an hour with correct dose, frequency and combo of pain meds. Prompt phone feedback to the vet is essential in getting pain relief meds properly adjusted for dose and frequency.	<ul> <li>2. SWELLING reduction         <ul> <li>in the spinal cord may take             anywhere from 7-30 days             before all painful swelling is             prooved absent.</li> </ul> </li> <li>Danger: Never give a         <ul> <li>NSAID with a steroid. 5-7             day washout needed with             switches:         <ul> <li>NSAID &lt;-&gt; NSAID</li> <li>Steroid &lt;-&gt; NSAID</li> </ul> </li> </ul></li></ul>	Takes 8 weeks during conservative treatment to self heal with 100% STRICT rest.	limit. Best to think in
MEDS: A combo to address the 3 sources of pain. Rx'd for every 8 hrs for round the clock control:	<b>MEDS:</b> There are two classes of drugs to bring down swelling in the spinal cord.	NO MEDS heal discs	NO MEDS heal nerves
<ul> <li>Tramadol- general pain relief</li> <li>Methocarbamol-muscle spasm pain</li> <li>Gabapentin-nerve pain</li> <li>Amantadine an extra dimension of relief for severe pain</li> <li>Signs of pain: <ul> <li>yelping</li> <li>trembling or shaking</li> <li>arched back</li> <li>reluctance to move</li> <li>tight/tense stomach</li> <li>holding a leg up and not wanting to use it</li> <li>dog's personality is not backnot interested in things</li> <li>head held in an unusual positionhead down or nose up high</li> <li>panting</li> </ul> </li> </ul>	<ul> <li>1. NSAID (non-steroid anti- inflammatory drug) <ul> <li>Rimadyl.</li> <li>Metacam, and others</li> </ul> </li> <li>2. Steroids <ul> <li>Prednisolone,</li> <li>Dexamethasone, and others</li> </ul> </li> <li>Stomach protection: <ul> <li>Pepcid AC (famotidine) suppresses stomach acids anti-inflamamtories cause in order to avoid bleeding ulcers.</li> </ul> </li> <li>Adverse signs needing prompt vet help: <ul> <li>Not eating or drinking</li> <li>Vomiting or w/blood</li> </ul> </li> <li>Diarrhea or w/blood</li> </ul> <li>2. Washout period of 5 to 7 days to protect the GI tract should there be a need to switch between NSAID brands or NSAID &lt;&gt; steroid. For emergencies w/o a washout, sucralfate is added to Pepcid AC for double GI tract protection.</li>	Movements to avoid with conservative treatment:: • baths • sleeping with you • sitting on your lap • sniff fests at potty time • Chiropractic therapy • water therapy • water therapy • eating outside the crate • dragging legs around Note: post-op dogs under the direction of the surgeon can begin PT, water therapy soon after surgery. PT is always deliberate, purposeful and under control. Post-op crate rest is usually for 6 weeks.	<ul> <li>Therapies::</li> <li>Laser light therapy</li> <li>Acupuncture</li> </ul> Acupuncture and Laser Therapy stimulate the cell's metabolism that leads to the body's natural repair abilities and can be started as soon as possible. NOTE: Laser light therapy contraindicated with tumors. Anytime out of the recovery suite is a dangerous time for the early healing disc during conservative treatment. Movement of the back/neck can increase a disc tear. For an animal with no or very mild neuro deficits, the risk of transporting to therapy has to be carefully weighed against what benefit is to be gained. Seek a mobile vet.